Checklist of the Birds of Jamaica



Please enter counts for every species observed and note any breeding evidence for entry into eBird Caribbean.

Black-bellied Whistling-Duck (Dendrocygna autumnalis)	Whimbrel (Numenius phaeopus)	Jamaican Owl* (Pseudoscops grammicus)
West Indian Whistling-Duck* (Dendrocygna arborea)	Ruddy Turnstone (Arenaria interpres)	Antillean Nighthawk (Chordeiles gundlachii)
American Wigeon (Anas americana)	Stilt Sandpiper (Calidris himantopus)	Northern Potoo* (Nyctibius jamaicensis)
Blue-winged Teal (Anas discors)	Sanderling (Calidris alba)	Black Swift (Cypseloides niger)
Northern Shoveler (Anas clypeata)	Least Sandpiper (Calidris minutilla)	White-collared Swift (Streptoprocne zonaris)
Ring-necked Duck (Aythya collaris)	White-rumped Sandpiper (Calidris fuscicollis)	Antillean Palm-Swift* (Tachornis phoenicobia)
Lesser Scaup (Aythya affinis)	Semipalmated Sandpiper (Calidris pusilla)	Jamaican Mango* (Anthracothorax mango)
Masked Duck* (Nomonyx dominicus)	Western Sandpiper (Calidris mauri)	Vervain Hummingbird* (Mellisuga minima)
Ruddy Duck* (Oxyura jamaicensis)	Short-billed Dowitcher (Limnodromus griseus)	Red-billed Streamertail* (Trochilus p. polytmus)
Least Grebe* (Tachybaptus dominicus)	Wilson's Snipe (Gallinago delicata)	Black-billed Streamertail* (Trochilus p. scitulus)
Pied-billed Grebe* (Podilymbus podiceps)	Spotted Sandpiper (Actitis macularius)	Jamaican Tody* (Todus todus)
White-tailed Tropicbird (Phaethon lepturus)	Solitary Sandpiper (<i>Tringa solitaria</i>)	Belted Kingfisher* (Megaceryle alcyon)
Magnificent Frigatebird* (Fregata magnificens)	Greater Yellowlegs (Tringa melanoleuca)	Jamaican Woodpecker* (Melanerpes radiolatus)
Brown Booby* (Sula leucogaster)	Willet* (Tringa semipalmata)	Yellow-bellied Sapsucker (Sphyrapicus varius)
Double-crested Cormorant) (Phalacrocorax auritus)	Lesser Yellowlegs (Tringa flavipes)	American Kestrel* (Falco sparverius)
Brown Pelican* (Pelecanus occidentalis)	Laughing Gull* (Leucophaeus atricilla)	Merlin (Falco columbarius)
Least Bittern* (Ixobrychus exilis)	Herring Gull (Larus argentatus)	Peregrine Falcon (Falco peregrinus)
Great Blue Heron* (Ardea herodias)	Lesser Black-backed Gull (Larus fuscus)	Black-billed Parrot* (Amazona agilis)
Great Egret* (Ardea alba)	Brown Noddy* (Anous stolidus)	Yellow-billed Parrot* (Amazona collaria)
Snowy Egret* (Egretta thula)	Bridled Tern (Onychoprion anaethetus)	Green-rumped Parrotlet* (Forpus passerinus)
Little Blue Heron* (Egretta caerulea)	Least Tern (Sternula antillarum)	Olive-throated Parakeet ^* (Eupsittula nana)
Tricolored Heron* (Egretta tricolor)	Gull-billed Tern (Gelochelidon nilotica)	Jamaican Elaenia* (Myiopagis cotta)
Reddish Egret* (Egretta rufescens)	Caspian Tern (Hydroprogne caspia)	Greater Antillean Elaenia* (Elaenia fallax)
Cattle Egret* (Bubulcus ibis)	Common Tern (Sterna hirundo)	Jamaican Pewee* (Contopus pallidus)
Green Heron* (Butorides virescens)	Forster's Tern (Sterna forsteri)	Sad Flycatcher* (Myiarchus barbirostris)
Black-crowned Night-Heron* (Nycticorax nycticorax)	Royal Tern* (Thalasseus maximus)	Rufous-tailed Flycatcher* (Myiarchus validus)
Yellow-crowned Night-Heron* (Nyctanassa violacea)	Sandwich Tern* (Thalasseus sandvicensis)	Stolid Flycatcher* (Myiarchus stolidus)
White Ibis* (Eudocimus albus)	Black Tern (Chlidonias niger)	Gray Kingbird* (Tyrannus dominicensis)
Glossy Ibis* (Plegadis falcinellus)	Rock Pigeon^* (Columba livia)	Loggerhead Kingbird* (Tyrannus caudifasciatus)
Turkey Vulture* (Cathartes aura)	White-crowned Pigeon* (Patagioenas leucocephala)	Jamaican Becard* (Pachyramphus niger)
Osprey* (Pandion haliaetus)	Plain Pigeon* (Patagioenas inornata)	Blue Mountain Vireo* (Vireo osburni)
Red-tailed Hawk* (Buteo jamaicensis)	Ring-tailed Pigeon* (Patagioenas caribaea)	Jamaican Vireo* (Vireo modestus)
Yellow-breasted Crake* (Hapalocrex flaviventer)	Eurasian Collared-Dove^* (Streptopelia decaocto)	Black-whiskered Vireo (Vireo altiloquus)
Clapper Rail* (Rallus crepitans)	Common Ground-Dove* (Columbina passerina)	Jamaican Crow* (Corvus jamaicensis)
Sora (Porzana Carolina)	Crested Quail-Dove* (Geotrygon versicolor)	Northern Rough-winged Swallow (Stelgidopteryx
Spotted Rail* (Pardirallus maculatus)	Ruddy Quail-Dove* (Geotrygon montana)	serripennis)
Purple Gallinule* (Porphyrio martinicus)	Caribbean Dove* (Leptotila jamaicensis)	Caribbean Martin (Progne dominicensis)
Common Gallinule* (Gallinula galeata)	White-winged Dove* (Zenaida asiatica)	Tree Swallow (Tachycineta bicolor)
American Coot* (Fulica americana)	Zenaida Dove* (Zenaida aurita)	Bank Swallow (Riparia riparia)
Limpkin* (Aramus guarauna)	Mourning Dove* (Zenaida macroura)	Barn Swallow (Hirundo rustica)
Black-necked Stilt* (Himantopus mexicanus)	Smooth-billed Ani* (Crotophaga ani)	Cave Swallow* (Petrochelidon fulva)
Black-bellied Plover (Pluvialis squatarola)	Yellow-billed Cuckoo (Coccyzus americanus)	Rufous-throated Solitaire* (Myadestes genibarbis
Wilson's Plover* (Charadrius wilsonia)	Mangrove Cuckoo* (Coccyzus minor)	Swainson's Thrush (Catharus ustulatus)
Semipalmated Plover (Charadrius semipalmatus)	Chestnut-bellied Cuckoo* (Coccyzus pluvialis)	White-eyed Thrush* (Turdus jamaicensis)
Killdeer* (Charadrius vociferus)	Jamaican Lizard-Cuckoo* (Coccyzus vetula)	White-chinned Thrush* (Turdus aurantius)
Northern Jacana* (Jacana spinose)	Barn Owl* (Tyto alba)	Gray Catbird (Dumetella carolinensis)

Bahama Mockingbird*(Mimus gundlachii)
 Northern Mockingbird* (Mimus polyglottos)
 European Starling^* (Sturnus vulgaris)
 Ovenbird (Seiurus aurocapilla)
 Worm-eating Warbler (Helmitheros vermivorum)
Louisiana Waterthrush (Parkesia motacilla)
Northern Waterthrush (Parkesia noveboracensis)
Black-and-white Warbler (Mniotilta varia)
Prothonotary Warbler (Protonotaria citrea)
 Swainson's Warbler (Limnothlypis swainsonii)
Common Yellowthroat (Geothlypis trichas)
Arrowhead Warbler* (Setophaga pharetra)
Hooded Warbler (Setophaga citrina)
_ American Redstart (Setophaga ruticilla)
_ Cape May Warbler (Setophaga tigrina)
Northern Parula (Setophaga americana)
Magnolia Warbler (Setophaga magnolia)
Blackburnian Warbler (Setophaga fusca)
Yellow Warbler* (Setophaga petechia)
Chestnut-sided Warbler (Setophaga pensylvanica)
 Blackpoll Warbler (Setophaga striata)
 Black-throated Blue Warbler (Setophaga caerulescens)
Palm Warbler (Setophaga palmarum)
_Yellow-rumped Warbler (Setophaga coronata)
Yellow-throated Warbler (Setophaga dominica)
Prairie Warbler (Setophaga discolor)
 Black-throated Green Warbler (Setophaga virens)
Saffron Finch^* (Sicalis flaveola)
Bananaquit* (Coereba flaveola)
Yellow-faced Grassquit* (Tiaris olivaceus)
Black-faced Grassquit* (Tiaris bicolor)
Orangequit* (Euneornis campestris)
 Greater Antillean Bullfinch* (Loxigilla violacea)
 Yellow-shouldered Grassquit* (Loxipasser anoxanthus)
Jamaican Spindalis* (Spindalis nigricephala)
Grasshopper Sparrow* (Ammodramus savannarum)
Summer Tanager (Piranga rubra)
Rose-breasted Grosbeak (Pheucticus Iudovicianus)
 Indigo Bunting (Passerina cyanea)
 _Jamaican Blackbird* (Nesopsar nigerrimus)
 Great-tailed Grackle* (Quiscalus mexicanus)
Greater Antillean Grackle* (Quiscalus niger)
Shiny Cowbird* (Molothrus bonariensis)
Jamaican Oriole* (Icterus leucopteryx)
Baltimore Oriole (Icterus galbula)
_Jamaican Euphonia* (Euphonia jamaica)
House Sparrow^* (Passer domesticus)
_Scaly-breasted Munia^* (Lonchura punctulata)
_ Tricolored Munia^* (Lonchura malacca)
Chestnut Munia^* (Lonchura atricanilla)

Red – Endemic species, Jamaica Green – Endemic species, West Indies ^ – Introduced species

* – Year-round resident

Note: This is a list of the birds most likely to be seen in Jamaica. It does not include all rarely observed species.

Additional Species:	
	Count Information
Location	
Observation date _	
Start time	Duration
Number of people i	n group
Count type (see def	finitions below)
We encourage you	to record all birds seen and heard.
heard? ☐ YES ☐ I	complete checklist of the birds seen and NO

Submitting Your Data

After you have completed your bird count, submit your data to eBird Caribbean at http://ebird.org/caribbean

Defining Your Count

Stationary Count: A count made from a fixed location (moving less than 30 m). Examples are counting birds from an observation tower or in your backyard.

Traveling Count: Observations made while birding over a specific distance (preferably less than 5 km) and duration. Examples include walking a trail, driving an auto loop, and birding on a boat trip. If you walk 2 km. on a trail and return via the same route, record the distance as 2 km. Only record new birds that you are certain you did not previously detect on the return. Note: Counts carried out at wetlands and beaches should be entered as Caribean Waterbird Census (CWC) counts.

Checklist of the Birds of Jamaica









Sponsored by eBird Caribbean